

# INTERNATIONAL TRAUMA LIFE SUPPORT

## ROLE OF TXA IN MANAGEMENT OF TRAUMATIC HEMORRHAGE IN THE FIELD

Roy Alson, MD, PhD, FACEP, FAAEM and Sabina Braithwaite, MD, MPH, FACEP

The guidelines and references contained in this document are current as of the date of publication and in no way replace physician medical oversight.

### INTRODUCTION

Hemorrhagic shock remains a serious problem for the multiple trauma patient. It is the leading cause of preventable trauma deaths after loss of airway. Rapid and effective control of exsanguinating hemorrhage has been demonstrated to markedly improve survival and outcome, especially in the combat situation. Increased use of tourniquets by combat medics has reduced death from hemorrhagic shock in the most recent wars fought by U.S. and NATO forces.

The tactical and military environment is associated with a higher percentage of penetrating trauma and external hemorrhage than is seen with the civilian sector, in which blunt trauma predominates. This leads to the situation of ongoing hemorrhage that is difficult to control. Prompt recognition of this with transport to the appropriate facility (Trauma Center) and limiting fluid resuscitation to the level of restoring perfusion (hypotensive resuscitation) have been shown to result in improved survival for the trauma patient.

### BACKGROUND

Tranexamic acid (TXA) is an antifibrinolytic that has been used for many years to assist with the management of spontaneous hemorrhaging in the hemophilia patient. The use of this agent for management of hemorrhage in combat wounds has been reported in several papers. One of the most significant findings in the CRASH-2 study is that the use of TXA is associated with a 1.5 absolute risk reduction for death from hemorrhage. Other studies show that TXA is most effective if given within 3 hours of the injury and may be detrimental if given after that time.



## CONSIDERATIONS

The side effects of the agent are minimal and the contraindications are few. It is administered as a simple IV infusion, does not require refrigeration or extensive laboratory studies to allow administration (as is seen with blood products) and is inexpensive. (NOTE: Use for traumatic hemorrhage is an off label use per FDA in the United States.)

## PROCEDURE

Based on local protocols and clearance, TXA should be considered in those patients who show signs of hemorrhagic shock, including tachycardia (>110 BPM) and hypotension (SBP<100) and are less than three hours from injury. Do not give TXA through the same line as blood products.

## MEDICAL OVERSIGHT

Medical oversight should review current literature and develop pre-hospital EMS protocols in regard to appropriate use of TXA. Implementation of this protocol should be monitored and supervised through a quality assurance program.

## CONCLUSION

ITLS believes that there is sufficient evidence to support the use of TXA in the management of traumatic hemorrhage in the adult patient, pursuant to system medical control approval. Following initial resuscitation including control of external bleeding and stabilization of airway, consideration should be given to administration of TXA during early stages of transport.

## REFERENCES

1. Morrison JJ, et al. Military application of tranexamic acid in trauma emergency resuscitation (MATTERs) study. Arch Surg, 2012 Feb; 147(2): 113–9.
2. Shakur H et al, Effects of tranexamic acid on death, vascular occlusive events, and blood transfusion in trauma patients with significant hemorrhage (CRASH-2): a randomized, placebo-controlled trial. Lancet, 2010 Jul 3; 376(9734): 23–32.
3. Kobayashi L, Costantini TW, Coimbra R. Hypovolemic shock resuscitation. The Surgical clinics of North America. 2012;92(6):1403-23.
4. Rappold JF, Pusateri AE. Tranexamic acid in remote damage control resuscitation. Transfusion. 2013;53 Suppl 1:96S-9S.



5. Collaborators C-. Effect of tranexamic acid in traumatic brain injury: a nested randomised, placebo controlled trial (CRASH-2 Intracranial Bleeding Study). *British Medical Journal*. 2011;343:d3795.
6. Ker K, Edwards P, Perel P, Shakur H, Roberts I. Effect of tranexamic acid on surgical bleeding: systematic review and cumulative meta-analysis. *Bmj*. 2012;344:e3054.
7. Lockey DJ, Weaver AE, Davies GE. Practical translation of hemorrhage control techniques to the civilian trauma scene. *Transfusion*. 2013;53 Suppl 1:17S-22S.
8. Cap AP, Baer DG, Orman JA, Aden J, Ryan K, Blackbourne LH. Tranexamic Acid for Trauma Patients: A Critical Review of the Literature. *Journal of Trauma-Injury Infection & Critical Care*. 2011;71(1 Supplemental):S9-S14.



*Improving Trauma Care Worldwide*

# *Current Thinking*

## **Role of TXA In Management of Traumatic Hemorrhage In The Field** International Trauma Life Support

The guidelines and references contained in this document are current as of the date of publication and in no way replace physician medical oversight.

### **Abstract**

This is the official current thinking of International Trauma Life Support (ITLS) with regard to the role of TXA in management of traumatic hemorrhage in the pre-hospital setting.

### **Current Thinking**

It is the position of International Trauma Life Support that:

1. There is sufficient evidence to support the use of TXA in the management of traumatic hemorrhage in adult trauma patients.
2. ITLS supports the use of TXA in the acute management of traumatic hemorrhagic shock within the framework of established system medical oversight and protocols.
3. Use of TXA is recommended in conjunction with initial resuscitation and control of external bleeding. Early TXA administration should be considered following airway stabilization, control of external bleeding, and initial volume resuscitation.
4. The use of TXA should be considered during the early stages of resuscitation and transport. Current research demonstrates TXA is most effective if given within 3 hours of the injury and may be detrimental if given after that time.

